

When $\text{Aut}(\mathcal{A})$ and $\text{Homeo}(\text{Prim}(\mathcal{A}))$ are homeomorphic, where $\mathcal{A}$ is a C^* -algebra

B. BOUALI

Abstract

In this paper, we discuss when $\text{Aut}(\mathcal{A})$ and $\text{Homeo}(\text{Prim}(\mathcal{A}))$ are homeomorphic, where $\mathcal{A}$ is a C^* -algebra.

A. M.S 2000 subject classification : 46H05

Keywords: Automorphism of C^* -algebras, Primitive ideals, Hull-kernel topology.

1 Preliminaries

Let $\mathcal{A}$ be C^* -algebra. Then the collection $\text{Aut}(\mathcal{A})$ of $\star$ -automorphisms of $\mathcal{A}$ is a group under composition. We give $\text{Aut}(\mathcal{A})$ the point-norm topology, that is, $\alpha_n \mapsto \alpha$ if and only if $\alpha_n(a) \mapsto \alpha(a)$ for all $a \in \mathcal{A}$. Then $\text{Aut}(\mathcal{A})$ is a topological group.

Let $(X, \mathcal{X})$ and $(Y, \mathcal{Y})$ be topological spaces, and let $\phi : X \mapsto Y$ be a map. Then ϕ is called continuous (from $(X, \mathcal{X})$ to $(Y, \mathcal{Y})$) if the preimage $\phi^{-1}(V)$ of V under ϕ belongs to $\mathcal{X}$ for each $V \in \mathcal{Y}$. A bijection between topological spaces is called a homeomorphism if it is continuous and has a continuous inverse.

1.1 The hull kernel topology

The topology on $\text{Prim}(\mathcal{A})$ is given by means of a closure operation. Given any subset W of $\text{Prim}(\mathcal{A})$, the closure $\overline{W}$ of W is by definition the set of all elements in $\text{Prim}(\mathcal{A})$ containing $\cap W = \{\cap I : I \in W\}$, namely

$$\overline{W} = \{I \in \text{Prim}(\mathcal{A}) : I \supseteq \cap W\}$$

It follows that the closure operation defines a topology on $\text{Prim}(\mathcal{A})$ which is called Jacobson topology or hull kernel topology [1].

Proposition 1.1: The space $\text{Prim}(\mathcal{A})$ is a T_0 -space, i.e. for any two distinct points of the space there is an open neighborhood of one of the points which does not contain the other.

Proposition 1.2: If $\mathcal{A}$ is a C^* -algebra, then $\text{Prim}(\mathcal{A})$ is locally compact. If $\mathcal{A}$ has a unit, then $\text{Prim}(\mathcal{A})$ is locally compact.

Remark 1.1: The set of $\mathcal{K}(\mathcal{H})$ of all compact operators on the Hilbert space $\mathcal{H}$ is the largest two sided ideal in the C^* -algebra $\mathcal{B}(\mathcal{H})$ of all bounded operators.

Definition 1.1: A C^* -algebra $\mathcal{A}$ is said to be liminal if for every irreducible representation $(\pi, \mathcal{H})$ of $\mathcal{A}$ one has that $\pi(\mathcal{A}) = \mathcal{K}(\mathcal{H})$.

So, the algebra $\mathcal{A}$ is liminal if it is mapped to the algebra of compact operators under any irreducible representation. Furthermore if $\mathcal{A}$ is a liminal algebra, then one can prove that each primitive ideal of $\mathcal{A}$ is automatically a maximal closed two-sided ideal. As a consequence, all points of $\text{Prim}(\mathcal{A})$ are closed and $\text{Prim}(\mathcal{A})$ is a T_1 -space. In particular, every commutative C^* -algebra is liminal [1].

2 $\text{Aut}(\mathcal{A})$ and $\text{Homeo}(\text{Prim}(\mathcal{A}))$

Theorem 2.1: Let $\mathcal{A}$ be a liminal C^* -algebra, If $\alpha \in \text{Aut}(\mathcal{A})$, there is $h \in \text{Homeo}(\text{Prim}(\mathcal{A}))$ such that :

$$\alpha(a)(\pi) = h^{-1}(\pi)(a) \quad \forall a \in \mathcal{A} \text{ and } \text{Ker}(\pi) \in \text{Prim}(\mathcal{A})$$

The map $\alpha \mapsto h$ is a homeomorphism.

Proof The primitive ideal of $\mathcal{A}$ is a maximal ideal [1, corollary 4.1.11.(ii)], Let I_π be a maximal ideal of $\mathcal{A}$ for some $\pi \in \mathcal{A}$. Since $\alpha^{-1}(I_\pi)$ is a maximal ideal, there is a function h

$$h : \text{Prim}(\mathcal{A}) \mapsto \text{Prim}(\mathcal{A})$$

such that $\alpha^{-1}(I_\pi) = I_{h(\pi)}$.

Since we can replace α^{-1} by α , it follows that h is a bijection. we have induced isomorphism $\chi_\pi : \mathcal{A}/I_\pi \mapsto \mathbf{C}$ given by $\chi_\pi(a) = \pi(a)$ and $\beta : \mathcal{A}/I_\pi \mapsto \mathcal{A}/I_{h^{-1}(\pi)}$ given by $\beta(a + I_\pi) = \alpha(a) + I_{h^{-1}(\pi)}$. Therefore we get a commutative diagram:

$$\begin{array}{ccc}
\mathcal{A}/I_\pi & \xrightarrow{\beta} & \mathcal{A}/I_{h(\pi)} \\
\chi_\pi \downarrow & & \downarrow \chi_{h(\pi)} \\
\mathbf{C} & \xrightarrow{\gamma} & \mathbf{C}
\end{array}$$

and an induced isomorphism $\gamma : \mathbf{C} \mapsto \mathbf{C}$ defined by

$$\gamma(\pi(a)) = h^{-1}(\pi)(\alpha(a))$$

Since γ must be the identity maps, then $\pi(a) = h^{-1}(\pi)(\alpha(a))$.

All opens set of $\text{Prim}(\mathcal{A})$ are of the forms $U_I = \{P \in \text{Prim}(\mathcal{A}) : P \not\supseteq I\}$ [2,3], calculate $h^{-1}(U_I)$.

$$\begin{aligned}
h^{-1}(U_I) &= \{ \pi : \ker(\pi) \in \text{Prim}(\mathcal{A}) \text{ and } \ker(\pi) \not\supseteq I \} \\
&= \{ h^{-1}(\pi) : \ker(\pi) \in \text{Prim}(\mathcal{A}) \text{ and } \ker(\pi) \not\supseteq I \} \\
&= \{ \pi' : \ker(\pi') \in \text{Prim}(\mathcal{A}) \text{ and } \ker(h(\pi')) \not\supseteq I \} \\
&= \{ \pi' : \ker(\pi') \in \text{Prim}(\mathcal{A}) \text{ and } \alpha^{-1}(I_\pi) \not\supseteq I \} \\
&= \{ \pi' : \ker(\pi') \in \text{Prim}(\mathcal{A}) \text{ and } \ker(\pi') \not\supseteq \alpha(I) \} \\
&= U_{\alpha(I)}.
\end{aligned}$$

Then $h^{-1}(U_I)$ is an open set and h is continuous. Replace α by α^{-1} , it follows that h^{-1} is continuous, then h is a homeomorphism.

We still have to check that $\alpha \mapsto h$ is a homeomorphism. Let α_n in $\text{Aut}(\mathcal{A})$ be given by $\pi(\alpha_n(a)) = h_n(\pi)(a)$.

Suppose that $\alpha_n \mapsto \alpha$. If, contrary there exists $\pi \in \text{Homeo}(\text{Prim}(\mathcal{A}))$ such that $h_n(\pi) \not\mapsto h(\pi)$ and $h_n(\pi)(a) \not\mapsto h(\pi)(a)$, then $\pi(\alpha_n(a)) \not\mapsto \pi(\alpha(a))$. Thus $\alpha_n(a) \mapsto \alpha(a)$ in $\mathcal{A}$. This is a contradiction, and we must have $h_n \mapsto h$ in $\text{Homeo}(\text{Prim}(\mathcal{A}))$. Since $\alpha_n^{-1} \mapsto \alpha^{-1}$, we have also $h_n^{-1} \mapsto h^{-1}$. Thus $h_n \mapsto h$ in $\text{Homeo}(\text{Prim}(\mathcal{A}))$.

Now suppose that $h_n \mapsto h$ in $\text{Homeo}(\text{Prim}(\mathcal{A}))$. We need to show that given $a \in \mathcal{A}$, and I_π in $\text{Prim}(\mathcal{A})$. If not, there exists $a \in \mathcal{A}$, such that $\alpha_n(a) \not\mapsto \alpha(a)$. Thus $h_n(\pi)(a) \not\mapsto h(\pi)(a)$. This is a contradiction, and we must have $\alpha_n \mapsto \alpha$ in $\text{Aut}(\mathcal{A})$.

References

- [1] J. Dixmier *Les C^* -algebres et leurs représentations*, Gauthier-villars éditeur, Paris 1969.

- [2] J. Dixmier *C*-algebras*, North-Holland, New York, 1977.